



Europäisch s Patentamt  
European Patent Office  
Office uropéen des brevets

THE BRITISH LIBRARY  
SCIENCE REFERENCE AND INFORMATION SERVICE

Publication number:

**0 253 500  
A2**

## EUROPEAN PATENT APPLICATION

Application number: 87305243.5

Int. Cl.4: C07C 103/76 , C07C 103/78

Date of filing: 12.06.87

Priority: 18.07.86 GB 8617652

Date of publication of application:  
20.01.88 Bulletin 88/03

Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE


Applicant: **IMPERIAL CHEMICAL INDUSTRIES  
PLC**  
Imperial Chemical House Millbank  
London SW1P 3JF(GB)

Inventor: **Tucker, Howard**  
32 Millers Meadow Rainow  
Macclesfield Cheshire(GB)

Representative: **Slatcher, Reginald Peter et al**  
Imperial Chemical Industries PLC Legal  
Department Patents P.O. Box 6 Bessemer  
Road  
Welwyn Garden City AL7 1HD(GB)

Acylanilide derivatives.

An acylanilide of the formula:



wherein R<sup>1</sup> or R<sup>2</sup> which may be the same or different, each is an electron-withdrawing substituent, alkylthio or phenylthio or R<sup>1</sup> is hydrogen, alkyl or alkoxy;  
wherein R<sup>3</sup> is hydrogen or halogen;  
wherein R<sup>4</sup> is hydrogen or acyl  
wherein A is branched-chain alkylene;  
and wherein R<sup>5</sup> is phenyl, substituted phenyl or naphthyl. The compounds possess progestational activity.


ACYLANILIDE DERIVATIVES

This invention relates to new acylanilide derivatives which possess progestational activity.

Many acylanilides chemically-related to those hereinafter described are known to possess antian-  
drogenic activity. Of these, flutamide and hydroxyflutamide have been known for some considerable time,  
and others are described in Applicant's European Specifications Nos. 2309, 2892, 40932, 79191 and  
100172.

We have now discovered that a small group of acylanilides surprisingly possess no antandrogenic  
activity, but potent progestational activity.

According to the invention there is provided an acylanilide of the formula:



wherein R<sup>1</sup> is cyano, carbamoyl, nitro, fluoro, chloro, bromo, iodo or hydrogen, or alkyl, alkoxy, alkanoyl  
alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or per-  
fluoroalkylsulphonyl each of up to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl;

wherein R<sup>2</sup> is cyano, carbamoyl, nitro, fluoro, chloro, bromo or iodo or alkanoyl, alkylthio, alkylsulphinyl,  
alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of up  
to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl;

wherein R<sup>3</sup> is hydrogen or halogen;

wherein R<sup>4</sup> is hydrogen or acyl of up to 15 carbon atoms;

wherein A is branched-chain alkylene of up to 6 carbon atoms;

and wherein R<sup>5</sup> is phenyl which bears one, two or three substituents selected from hydrogen, halogen, nitro,  
hydroxy, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl,  
perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkyl-  
carbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl; or  
R<sup>5</sup> is naphthyl.

It will be observed that the acylanilide derivative of the invention possesses an asymmetric carbon  
atom, namely the carbon atom which bears the substituent OR<sup>4</sup>, and it can therefore exist in racemic and  
optically-active forms. Furthermore, frequently the branched-chain alkylene group -A- will also possess an  
asymmetric carbon atom in which case pairs of diastereoisomers, in optically-active or racemic form, are  
also possible. It is to be understood that this invention encompasses any racemic form of the acylanilide  
derivative and any optically-active form which possesses progestational activity, it being a matter of  
common general knowledge how a racemic compound may be resolved into its optically-active forms and  
how any progestational activity present in any of these forms may be determined.

A suitable value for R<sup>1</sup> when it is alkyl, or for an alkyl substituent in R<sup>5</sup> when R<sup>5</sup> is phenyl substituted by  
alkyl is, for example, methyl or ethyl.

A suitable value for R<sup>1</sup> when it is alkoxy, or for an alkoxy substituent in R<sup>5</sup> when R<sup>5</sup> is phenyl substituted  
by alkoxy is, for example, methoxy or ethoxy.

A suitable value for R<sup>1</sup> or R<sup>2</sup> when it is alkanoyl, or for an alkanoyl substituent in R<sup>5</sup> when R<sup>5</sup> is phenyl  
substituted by alkanoyl is, for example, formyl or acetyl.

A suitable value for R<sup>1</sup> or R<sup>2</sup> when it is alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, per-  
fluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl, or for such a substituent in R<sup>5</sup> when R<sup>5</sup> is  
phenyl bearing such a substituent is, for example, methylthio, ethylthio, methylsulphinyl, methylsulphonyl,  
trifluoromethyl, pentafluoroethyl, trifluoromethylthio, trifluoromethylsulphinyl or trifluoromethylsulphonyl.

A suitable value for R<sup>3</sup> when it is halogen, or for a halogen substituent in R<sup>5</sup> when R<sup>5</sup> is phenyl  
substituted by halogen, is fluoro, chloro, bromo or iodo.

R<sup>3</sup> is preferably hydrogen or chloro, especially hydrogen.

A suitable value for an alkoxycarbonyl or N-alkylcarbamoyl substituent in R<sup>5</sup> when R<sup>5</sup> is phenyl bearing  
such a substituent is, for example, methoxycarbonyl, ethoxycarbonyl or N-methylcarbamoyl.

A suitable value for R<sup>4</sup> when it is acyl is, for example, alkanoyl or aroyl each of up to 15 carbon atoms, for example acetyl, propionyl, decanoyl, dodecanoyl or benzoyl.

R<sup>4</sup> is preferably hydrogen,

A suitable value for A is, for example, 2-methylethylene or 2,2-dimethylethylene.

A preferred combination of values for R<sup>1</sup> and R<sup>2</sup> is as follows:-

	R <sup>1</sup>	R <sup>2</sup>
10	trifluoromethyl	nitro
	trifluoromethyl	cyano
15	chloro	chloro
	chloro	nitro
	chloro	cyano
	cyano	cyano
20	nitro	cyano
	ethoxy	nitro
	chloro	methylsulphonyl
25		


A preferred acylanilide of the invention has the formula stated above wherein R<sup>1</sup> and R<sup>2</sup>, which may be the same or different, each is cyano, nitro, trifluoromethyl, methylthio, methylsulphinyl, methylsulphonyl or chloro, R<sup>3</sup> and R<sup>4</sup> are both hydrogen, A is 2-methylethylene or 2,2-dimethylethylene and R<sup>5</sup> is phenyl which is unsubstituted or which bears one fluoro, chloro, hydroxy, methyl, nitro, methylthio, methylsulphinyl or methylsulphonyl substituent.

Specific acylanilides of the invention are hereinafter described in the Examples.


A particularly active compound is N(2-hydroxy-4-phenyl-2-trifluoromethylpentanoyl)-4-nitro-3-trifluoromethylaniline in either of its diastereoisomeric forms.

The acylanilides of the invention may be manufactured by any chemical process known to be suitable for the manufacture of chemically-analogous compounds.

A preferred process for the manufacture of an acylanilide of the invention comprises the reaction of an amine of the formula:-



wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> have the meanings stated above, with an acid of the formula:-



10 wherein  $\text{R}^4$ ,  $\text{R}^5$  and A have the meanings stated above, or with a reactive derivative of said acid.

A suitable reactive derivative of an acid is, for example, an acid anhydride, or an acyl halide, for example an acyl chloride, or a lower alkyl ester of said acid, for example the methyl or ethyl ester.

Preferably the reaction is carried out in N,N-dimethylacetamide solution using an acyl chloride (prepared from the acid and thionyl chloride) as reactant.

15 An acylanilide of the invention wherein  $\text{R}^4$  is hydrogen may be prepared by the hydrolysis of the corresponding acylanilide wherein  $\text{R}^4$  is acyl, and conversely, an acylanilide of the invention wherein  $\text{R}^4$  is acyl may be prepared by the acylation of the corresponding acylanilide wherein  $\text{R}^4$  is hydrogen.

An acylanilide of the invention wherein one or more of  $\text{R}^1$ ,  $\text{R}^2$  and a substituent in the phenyl group  $\text{R}^5$  is alkylsulphinyl, perfluoroalkylsulphinyl or phenylsulphinyl, or is alkylsulphonyl, perfluoroalkylsulphonyl or phenylsulphonyl, may be prepared by the oxidation of the corresponding acylanilide wherein one or more of  $\text{R}^1$ ,  $\text{R}^2$  and a substituent in the phenyl group  $\text{R}^5$  is alkylthio, perfluoroalkylthio or phenylthio, respectively. The oxidising agent and conditions used will determine whether a sulphinyl or a sulphonyl compound is obtained. Thus, oxidation with sodium metaperiodate in methanol solution at or below laboratory temperature will generally convert a thio compound into the corresponding sulphinyl compound; and oxidation with a peracid, for example m-chloroperbenzoic acid in methylene chloride solution at or above laboratory temperature will generally convert a thio compound into the corresponding sulphonyl compound.

25 A racemic acylanilide of the invention wherein  $\text{R}^4$  is hydrogen may be separated into its optical isomers by forming an ester of the hydroxy group  $\text{OR}^4$  with an optically-active acid, for example (-)-camphanic acid, separating the diastereoisomeric esters thus obtained, by fractional crystallisation or, preferably, by flash-chromatography, and then hydrolysis of each separate ester to the alcohol.

As stated above, an acylanilide of the invention possesses progestational properties as demonstrated by its ability to promote glandular development in the endometrium of an oestrogen-primed immature rabbit, the standard Clauberg assay procedure. An acylanilide of the invention may therefore be used as an oral contraceptive, and in the treatment of menstrual disorders, such as dysmenorrhea, dysfunctional bleeding and premenstrual tension, and in the treatment of hormone dependent tumors, especially those of the breast or endometrium. It may also be used for the synchronisation of oestrus and for the maintenance of early pregnancy in domestic animals such as cattle. At a dose of an acylanilide of the invention which produces progestational activity in rabbits no symptoms of toxicity are apparent.

35 The acylanilide of the invention may be administered to a warm blood animal in the form of a pharmaceutical or veterinary composition which comprises the acylanilide in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral dosage, as a tablet, capsule, aqueous or oily solution or suspension or emulsion. It may alternatively be in the form of a sterile solution or suspension suitable for parenteral administration, or be in the form of an ointment or lotion for topical administration or be in the form of a suppository for anal or vaginal administration.

45 The composition may additionally contain one or more drugs selected from oestrogens, for example, ethnyloestradiol or mestranol (which combination may be used as an oral contraceptive); antioestrogens, for example tamoxifen; androgens, for example cyproterone acetate and methyltestosterone; and gonadotrophin releasing factors and analogues thereof and antagonists thereof.

50 The acylanilide of the invention will normally be administered to a warm-blooded animal at a dose of between 0.1 mg. and 125 mg. per kg. bodyweight.

The invention is illustrated but not limited by the following Examples:-

Example 1

Thionyl chloride (0.82 ml.) was added dropwise to a stirred solution of 2-hydroxy-4-phenyl-2-trifluoromethylpentanoic acid (Isomer A, m.p. 145-146°C.) in dimethylacetamide (60 ml.) which was maintained at -15°C. and under an atmosphere of argon, the mixture was stirred at that temperature for 1 hour and 4-nitro-3-trifluoromethylaniline (1.9 g.) was added. The mixture was stirred at -15°C. for 1 hour and then at laboratory temperature for 20 hours, and was then poured into water (700 ml.). The mixture was extracted six times with diethyl ether (100 ml.) each time) and the combined extracts were washed twice with aqueous 10% w/v sodium carbonate solution (50 ml. each time), dried over magnesium sulphate and evaporated to dryness. The residue was purified by flash chromatography on a silica gel column (Merck 9835) using a 2:1 v/v mixture of petroleum ether (b.p. 60-80°C.) and ethyl acetate as eluant. There was thus obtained N-(2-hydroxy-4-phenyl-2-trifluoromethylpentanoyl)-4-nitro-3-trifluoromethylaniline, m.p. 162-163°C. (more polar isomer).

The pentanoic acid used as starting material was obtained as follows:-

1,1,1-Trifluoromethyl-4-phenylpentan-2-one (116.7 g.; b.p. 82-83°C./10 mm. Hg.; prepared by the general process described in the Journal of Organic Chemistry, 1967, 32, 1316) was added dropwise to a cooled stirred solution of potassium cyanide (37.85 g.) in water (240 ml.) at such a rate that the temperature of the mixture was maintained at between 0°C. and 5°C. A 4:1 v/v mixture of water and concentrated sulphuric acid (300 ml.) was added at such a rate as to maintain the above temperature, and the mixture was stirred at laboratory temperature for 15 hours and then extracted four times with diethyl ether (90 ml. each time). The combined extracts were washed three times with water (100 ml. each time), dried over magnesium sulphate and evaporated to dryness under reduced pressure.


A mixture of the cyanhydrin thus obtained as a mixture of diastereoisomers (6.0 g.), concentrated aqueous hydrochloric acid (45 ml.) and glacial acetic acid (15 ml.) was vigorously stirred and heated at 100°C. for 48 hours, cooled and poured onto ice (125 g.). The mixture was extracted three times with diethyl ether (50 ml. each time) and the combined extracts were washed with water (50 ml.) and then extracted three times with saturated aqueous sodium carbonate solution (50 ml. each time). The combined extracts were acidified with concentrated aqueous hydrochloric acid and extracted three times with diethyl ether (50 ml. each time). The combined extracts were washed with water (50 ml.) and saturated aqueous sodium chloride solution (50 ml.), dried over magnesium sulphate and evaporated to dryness. The residue was stirred with petroleum ether (b.p. 60-80°C.) and the mixture was filtered. There was thus obtained as solid residue Isomer A of 2-hydroxy-4-phenyl-2-trifluoromethylpentanoic acid, m.p. 145-146°C. The petroleum ether filtrate was evaporated to dryness and there was thus obtained as residue a solid, m.p. 101-102°C., which was predominantly Isomer B of 2-hydroxy-4-phenyl-2-trifluoromethylpentanoic acid contaminated with a small amount of Isomer A.

Example 2

The process described in Example 1 was repeated using the impure Isomer B of the pentanoic acid (m.p. 101-102°C.) as starting material in place of Isomer A. The product was purified by flash chromatography on silica gel (Merck 9385) using a 3:2 v/v mixture of petroleum ether (b.p. 60-80°C.) and ethyl acetate as eluant, and the product obtained was crystallised from a 2:1 v/v mixture of ethyl acetate and petroleum ether (b.p. 60-80°C.). There was thus obtained N-(2-hydroxy-4-phenyl-2-trifluoromethylpentanoyl)-4-nitro-3-trifluoromethylaniline, m.p. 145-146°C. (less polar isomer).

## Claims

1. An acylanilide of the formula:



wherein R<sup>1</sup> is cyano, carbamoyl, nitro, fluoro, chloro, bromo, iodo or hydrogen, or alkyl, alkoxy, alkanoyl alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of up to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl; wherein R<sup>2</sup> is cyano, carbamoyl, nitro, fluoro, chloro, bromo or iodo or alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of up to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl;

wherein R<sup>3</sup> is hydrogen or halogen;

wherein R<sup>4</sup> is hydrogen or acyl of up to 15 carbon atoms;

wherein A is branched-chain alkylene of up to 6 carbon atoms;


and wherein R<sup>5</sup> is phenyl which bears one, two or three substituents selected from hydrogen, halogen, nitro, hydroxy, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl; or R<sup>5</sup> is naphthyl.

2. An acylanilide as claimed in claim 1, wherein R<sup>1</sup> and R<sup>2</sup>, which may be the same or different, each is cyano, nitro, trifluoromethyl, methylthio, methylsulphinyl, methylsulphonyl or chloro, R<sup>3</sup> and R<sup>4</sup> are both hydrogen, A is 2-methylethylene or 2,2-dimethylethylene and R<sup>5</sup> is phenyl which is unsubstituted or which bears one fluoro, chloro, hydroxy, methyl, nitro, methylthio, methylsulphinyl methylsulphonyl substituent.


3. The compound N-(2-hydroxy-4-phenyl-2-trifluoromethylpentanoyl)-4-nitro-3-trifluoromethylaniline in either of its diastereoisomeric forms.

4. A process for the manufacture of an acylanilide claimed in claim 1 which comprises:-

(a) the reaction of an amine of the formula:-



wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> have the meanings stated in claim 1 with an acid of the formula:-



wherein R<sup>4</sup>, R<sup>5</sup> and A have the meanings stated in claim 1 or with a reactive derivative of said acid;

(b) for the manufacture of an acylanilide of the invention wherein  $R^4$  is hydrogen, the hydrolysis of the corresponding anylanilide wherein  $R^4$  is acyl;

(c) for the manufacture of an acylanilide of the invention wherein  $R^4$  is acyl, the acylation of the corresponding acylanilide wherein  $R^4$  is hydrogen;

(d) for manufacture of an acylanilide of the invention wherein one or more of  $R^1$ ,  $R^2$  and a substituent in the phenyl group  $R^5$  is alkylsulphinyl, perfluoroalkylsulphinyl or phenylsulphinyl, or is alkylsulphonyl, perfluoroalkylsulphonyl or phenylsulphonyl, the oxidation of the corresponding acylanilide wherein one or more of  $R^1$ ,  $R^2$  and a substituent in the phenyl group  $R^5$  is alkylthio, perfluoroalkylthio or phenylthio, respectively or

(e) for the separation into its optical isomers of an acylanilide of the invention wherein  $R^4$  is hydrogen, the formation of an ester of the hydroxy group  $OR^4$  with an optically-active acid, the separation of the diastereoisomeric esters thus obtained by fractional crystallisation or by flash-chromatography, and then the hydrolysis of each separate ester to the alcohol.

5. A pharmaceutical or veterinary composition comprising an acylanilide as claimed in claim 1. in association with a pharmaceutically-acceptable diluent or carrier.

6. A composition as claimed in claim 5 which is in a form suitable for oral dosage, as a tablet, capsule, aqueous or oily solution or suspension or emulsion; or in the form of a sterile solution or suspension suitable for parenteral administration, or in the form of an ointment or lotion for topical administration or in the form of a suppository for anal or vaginal administration.


7. A composition as claimed in claim 5 which additionally contains one or more drugs selected from oestrogens, antioestrogens, androgens and gonadotrophin releasing factors and analogues thereof and antagonists thereof.

8. The use of an acylanilide as claimed in claim 1 for the manufacture of a medicament for producing a progestational effect in a warm-blooded animal.

9. The use of an acylanilide as claimed in claim 1 for producing a progestational effect in the human or animal body.

Claims for the following contracting states: Austria, Greece and Spain

1. A process for the manufacture of an acylanilide of the formula:-



wherein  $R^1$  is cyano, carbamoyl, nitro, fluoro, chloro, bromo, iodo or hydrogen, or alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of up to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl;

wherein  $R^2$  is cyano, carbamoyl, nitro, fluoro, chloro, bromo or iodo or alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of up to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl;

wherein  $R^3$  is hydrogen or halogen;


wherein  $R^4$  is hydrogen or acyl of up to 15 carbon atoms;

wherein A is branched-chain alkylene of up to 6 carbon atoms;


and wherein  $R^5$  is phenyl which bears one, two or three substituents selected from hydrogen, halogen, nitro, hydroxy, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl, or  $R^5$  is naphthyl characterised by:-

(a) the reaction of an amine of the formula:-

0 253 500



wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> have the meanings stated above, or with an acid of the formula:-



wherein R<sup>4</sup>, R<sup>5</sup> and A have the meanings stated above, or with a reactive derivative of said acid; or

(b) for the manufacture of an acylanilide of the invention wherein R<sup>4</sup> is hydrogen, the hydrolysis of the corresponding acylanilide wherein R<sup>4</sup> is acyl; or

(c) for the manufacture of an acylanilide of the invention wherein R<sup>4</sup> is acyl, the acylation of the corresponding acylanilide wherein R<sup>4</sup> is hydrogen; or

(d) for the manufacture of an acylanilide of the invention wherein one or more of R<sup>1</sup>, R<sup>2</sup> and a substituent in the phenyl group R<sup>5</sup> is alkylsulphinyl, perfluoroalkylsulphinyl or phenylsulphinyl, or is alkylsulphonyl, perfluoroalkylsulphonyl or phenylsulphonyl, the oxidation of the corresponding acylanilide wherein one or more of R<sup>1</sup>, R<sup>2</sup> and a substituent in the phenyl group R<sup>5</sup> is alkylthio, perfluoroalkylthio or phenylthio, respectively; or

(e) for the separation into its optical isomers of an acylanilide of the invention wherein R<sup>4</sup> is hydrogen, the formation of an ester of the hydroxy group OR<sup>4</sup> with an optically-active acid, the separation of the diastereoisomeric esters thus obtained by fractional crystallisation or by flash-chromatography, and then the hydrolysis of each separate ester to the alcohol.

2. A process as claimed in clause (a), (b), (d) or (e) of claim 1, wherein R<sup>1</sup> and R<sup>2</sup> which may be the same or different, each is cyano, nitro, trifluoromethyl, methylthio, methylsulphinyl, methylsulphonyl or chloro, R<sup>3</sup> and R<sup>4</sup> are both hydrogen, A is 2-methylethylene or 2,2-dimethylethylene and R<sup>5</sup> is phenyl which is unsubstituted or which bears one fluoro, chloro, hydroxy, methyl, nitro, methylthio, methylsulphinyl or methylsulphonyl substituent.

3. A process as claimed in clause (a), (b) or (e) of claim 1, wherein in the starting materials R<sup>1</sup> is trifluoromethyl, R<sup>2</sup> is nitro, R<sup>3</sup> is hydrogen, R<sup>4</sup> is hydrogen or acyl, A is 2-methylethylene and R<sup>5</sup> is phenyl, whereafter if R<sup>4</sup> is acyl the compound is hydrolysed to the corresponding compound wherein R<sup>4</sup> is hydrogen.



(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 87305243.5

(22) Date of filing: 12.06.87

(64) Int. Cl. **C07C 103/76**, **C07C 103/78**  
 THE BRITISH LIBRARY  
 SCIENCE REFERENCE AND INFORMATION SERVICE

(30) Priority: 18.07.86 GB 8617652

(43) Date of publication of application:  
20.01.88 Bulletin 88/03

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(88) Date of deferred publication of the search report:  
04.01.89 Bulletin 89/01


(71) Applicant: **IMPERIAL CHEMICAL INDUSTRIES PLC**  
**Imperial Chemical House Millbank**  
**London SW1P 3JF(GB)**

(72) Inventor: **Tucker, Howard**  
**32 Millers Meadow Rainow**  
**Macclesfield Cheshire(GB)**

(74) Representative: **Slatcher, Reginald Peter et al**  
**Imperial Chemical Industries PLC Legal**  
**Department: Patents P.O. Box 6 Bessemer**  
**Road**  
**Welwyn Garden City AL7 1HD(GB)**

(54) **Acylanilide derivatives.**

(57) An acylanilide of the formula:



wherein R<sup>1</sup> or R<sup>2</sup> which may be the same or different, each is an electron-withdrawing substituent, alkylthio or phenylthio or R<sup>1</sup> is hydrogen, alkyl or alkoxy;

wherein R<sup>3</sup> is hydrogen or halogen;

wherein R<sup>4</sup> is hydrogen or acyl

wherein A is branched-chain alkylene;

and wherein R<sup>5</sup> is phenyl, substituted phenyl or naphthyl. The compounds possess progestational activity.

**EP 0 253 500 A3**



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 87 30 5243

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A,D	EP-A-0 079 191 (ICI) * claims 1, 4-8 * ---	1-9	C 07 C 103/76 C 07 C 103/78 C 07 C 121/52 C 07 C 147/06 C 07 C 145/00 C 07 C 103/30
A,D	EP-A-0 100 172 (ICI) * claim 1 * -----	1,4	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 C 103/00 C 07 C 121/00 C 07 C 147/00 C 07 C 145/00
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 27-09-1988	Examiner RUFET J.M.A.
<div>CATEGORY OF CITED DOCUMENTS</div> <div>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</div> <div>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- &amp; : member of the same patent family, corresponding document</div>			